

STRONG BALMER LINES IN OLD STELLAR POPULATIONS: NO NEED FOR YOUNG AGES IN ELLIPTICALS?

CLAUDIA MARASTON AND DANIEL THOMAS

Universitäts-Sternwarte München, Scheinerstr. 1, D-81679 München, Germany

Accepted for publication in the Astrophysical Journal

ABSTRACT

Comparing models of Simple Stellar Populations (SSP) with observed line strengths generally provides a tool to break the age-metallicity degeneracy in elliptical galaxies. Due to the wide range of Balmer line strengths observed, ellipticals have been interpreted to exhibit an appreciable scatter in age. In this paper, we analyze Composite Stellar Population models with a simple mix of an old metal-rich and an old metal-poor component. We show that these models simultaneously produce strong Balmer lines and strong metallic lines without invoking a young population. The key to this result is that our models are based on SSPs that better match the steep increase of $H\beta$ in metal-poor globular clusters than models in the literature. Hence, the scatter of $H\beta$ observed in cluster and luminous field elliptical galaxies can be explained by a spread in the metallicity of *old* stellar populations. We check our model with respect to the so-called G-dwarf problem in ellipticals. For a galaxy subsample covering a large range in $1500-V$ colors we demonstrate that the addition of an old metal-poor subcomponent does not invalidate other observational constraints like colors and the flux in the mid-UV.

Subject headings: galaxies: elliptical and lenticular, cD – galaxies: stellar content – galaxies: formation – galaxies: abundances – galaxies: fundamental parameters (ages, metallicities)

1. INTRODUCTION

More than 20 years ago it has been recognized that the modeling of the spectral energy distribution of ellipticals is affected by an ambiguity in age and metallicity (Faber 1972; O’Connell 1976), which has turned out to be a general complication in population synthesis (Renzini 1986). However, considering Simple Stellar Populations (SSP) in the two-parameter space of Balmer and metallic lines, the age-metallicity degeneracy can be broken (González 1993; Worthey 1994). The major reason for the success of this strategy is that the Balmer line strengths of SSPs are predominantly age sensitive at metallicities above $\sim 1/3 Z_{\odot}$ that are supposed to be the only relevant for elliptical galaxies. Strong $H\beta$ lines are thus taken as an indication for young (intermediate-age) populations, the observed scatter is interpreted as a considerable spread in age (e.g., Faber et al. 1995).

An alternative path to obtain blue stars and hence strong Balmer lines is to consider old *metal-poor* populations. In this paper we follow this approach and compute composite stellar populations that contain an old metal-poor subcomponent. The principal focus is to check if a combination of only old populations can reproduce strong $H\beta$ without invalidating further constraints for ellipticals like metallic indices, colors and spectral energy distributions.

The paper is organized as follows. In Section 2 we calibrate the SSP model indices and spectral energy distributions on globular cluster data. The composite models and their application to galaxy data are presented in Section 3. In Sections 4 and 5 we discuss and summarize the results.

2. SIMPLE STELLAR POPULATION MODELS

We compute a new set of SSP models based on the population synthesis presented in Maraston (1998), in which the fuel consumption theorem (Renzini & Buzzoni 1986) is adopted to evaluate the energetics of the post main sequence phases. The new SSP models cover the metallicities $-2.25 < [\text{Fe}/\text{H}] < 0.5$

and ages $10^7 - 2 \cdot 10^{10}$ yr. The input stellar tracks are taken from Bono et al. (1997) and S. Cassisi (1999, private communication). These new models will be discussed in detail in a future paper (C. Maraston, in preparation). The calibration of the model colors on globular clusters is presented in Maraston (1998, 2000). Here we show the calibration of indices and spectral energy distributions of the SSP models that are relevant for the present study.

2.1. Spectral line indices

Synthetic line indices for SSPs are computed using the fitting functions from Worthey et al. (1994). Fig. 1 shows $H\beta$, Fe5335, and Mgb as a function of $[\text{Fe}/\text{H}]$ for galactic globular clusters from various data sets (see the caption). Our SSP models of fixed age (15 Gyr) and various metallicities are plotted as solid lines. Dotted lines are the models from Worthey (1994) for $t = 17$ Gyr.

As metal-poor stars ($[\text{Fe}/\text{H}] \lesssim -1.3$) spend their horizontal branch lifetimes at temperatures $T_{\text{eff}} \gtrsim 5500$ K, $H\beta$ is steeply rising with decreasing $[\text{Fe}/\text{H}]$ (upper panel in Fig. 1). This observed feature is the key for the present analysis and is well reproduced by our SSPs. The models from Worthey (1994), instead, predict a shallower trend, which implies that the very metal-poor globulars are younger than the more metal-rich ones. The synthetic Fe5335 indices are in good agreement with the Covino, Galletti & Pasinetti (1995) data, while for $[\text{Fe}/\text{H}] \gtrsim -1.5$ Burstein et al. (1984) and Trager (1998) derive lower values. Mgb indices seem slightly overestimated by the models at low metallicities.

Our synthetic metallic indices Fe5335, Mgb and the Balmer line $H\beta$ for $[\text{Fe}/\text{H}] \gtrsim -1.5$ agree well with the models from Worthey (1994), because both sets of models are based on the same fitting functions (Worthey et al. 1994). The uncertainty introduced by the use of different fitting functions is discussed in Maraston, Greggio & Thomas (1999) and L. Greggio & C. Maraston (in preparation). The low $H\beta$ values given by Worthey (1994) at very low metallicities are likely due to the




Fig. 1.— Calibration of the SSP models on galactic globular clusters. Data are from Burstein et al. 1984 (squares), Covino, Galletti & Pasinetti 1995 (circles), and Trager 1998 (triangles). $[\text{Fe}/\text{H}]$ (Zinn & West (1984) scale) is taken from Harris (1996). Solid lines are our SSPs for age $t = 15$ Gyr. Worthey (1994) SSPs for $t = 17$ Gyr are shown as dotted lines.

assumption of too red horizontal branch morphologies in the models.

As discussed in Worthey et al. (1994), iron indices are favorable as metallicity indicators, because the analysis with Mg-indices is severely affected by α -enhancement (Tripicco & Bell 1995): SSP models based on solar abundance ratios predict – for a given Fe index – weaker Mg indices than measured in elliptical galaxies (Worthey, Faber & González 1992; Greggio 1997). As at high metallicities the fitting functions of Fe5270 are significantly more uncertain (Maraston et al. 1999), in the following analysis we focus on $H\beta$ and Fe5335. We additionally consider the color $B-K$ as metallicity indicator which turns out to be in good agreement with Fe5335, while Mg b systematically implies higher metallicities (see Section 3).

Combining $H\beta$ with Fe5335, for 47 Tuc we obtain a ‘spectroscopic age’ of 15 Gyr (Fig. 1) in agreement with the age derived from the color-magnitude diagram (14 ± 1 Gyr, Richer et al. 1996). This example shows that the $H\beta$ -Fe5335 plane is successful in breaking the age-metallicity degeneracy, when applied to simple systems like globular clusters. Note that the

high resolution Balmer index $H\gamma_{\text{HR}}$, instead, leads to a ‘spectroscopic age’ in excess of 20 Gyr (Gibson et al. 2000).

2.2. Spectral energy distributions

The evolutionary synthesis code of Maraston (1998) is updated for the computation of the SSP spectral energy distributions (SEDs). The spectral library of Lejeune, Cuisinier & Buser (1998) is adopted to describe the stellar spectra as functions of gravity, temperature and metallicity.

The tightest constraint on the amount and the metallicity of metal-poor stars in elliptical galaxies comes from the flux in the mid-UV (2000–4000 Å). In Fig. 2 we present the calibration of synthetic metal-poor SEDs on galactic globular cluster data in the wavelength range $\lambda = 1500 - 3300$ Å (van Albada, de Boer & Dickens 1981). The model spectra are in excellent agreement with the data in the whole metallicity range. The increase of the flux shortward 3000 Å with decreasing metallicity is due to a hotter main sequence turn-off and a bluer horizontal branch. Note, however, that there are globulars of intermediate metallicity ($[\text{Fe}/\text{H}] \sim -1.5$) that show horizontal branches bluer




Fig. 2.— Calibration of the synthetic SEDs in the UV on galactic globular clusters. Data (squares) are taken from van Albada, de Boer & Dickens (1981), $[\text{Fe}/\text{H}]$ (Zinn & West (1984) scale) is adopted from Harris (1996). The filled squares (central panel) show M13 that has an extreme blue horizontal branch morphology. The solid lines are our model SEDs ($t = 15$ Gyr) with metallicities as indicated in the panels.

than what it is expected with the canonical mass-loss on the Red Giant Branch (van Albada et al. 1981). This case is shown in the central panel of Fig. 2 for the globulars M13 (NGC 6205), filled squares) and NGC 5272, which have the same metallicity $[\text{Fe}/\text{H}] \sim -1.54$ but different horizontal branch morphologies. M13 contains extreme blue horizontal branch stars (‘blue tails’), and exhibits an excess of flux shortward ~ 2000 Å (see Section 3.2.1).

As blue tails are preferentially found in denser and more concentrated globulars, it is likely that the particular dynamical conditions in such ‘second parameter’ globular clusters lead to enhanced mass-loss and extremely hot horizontal branch stars (Fusi Pecci et al. 1993), which may not apply to the average stellar field population in galaxies. Since in early-type galaxies globular clusters contribute less than 0.6 per cent to the total V-light (Ashman & Zepf 1998), we adopt the canonical mass loss $\eta = 0.33$ for $[\text{Fe}/\text{H}] \sim -1.5$ calibrated on ‘normal’ globulars like NGC 5272 (Fusi Pecci & Renzini 1975).

3. RESULTS

In the following we compare old (> 12 Gyr) composite stellar populations with data of elliptical galaxies. The models consist of two components: a major metal-rich population ($Z = Z_{\text{maj}}$) and a metal-poor subpopulation ($Z = Z_{\text{sub}}$). More complex models with a continuum of metallicities will be subject of a future paper.

3.1. Indices and colors

Fig. 3 displays $\text{H}\beta$ vs. $\text{Fe}5335$ (left-hand panel) and $\text{H}\beta$ vs. $B-K$ (right-hand panel) for composite models in which the ages of the major population and of the metal-poor subcomponent are fixed to 14 Gyr and 15 Gyr, respectively. The contribution from the low-metallicity population is 10 per cent by mass.

The grid in Fig. 3 shows models for various metallicities of the two components. Solid lines connect models with constant metallicity of the major component $Z_{\text{maj}} = 0.5, 1, 2, 3 Z_{\odot}$ (from left to right). The metallicity of the subcomponent is $Z_{\text{sub}} = 0.05$

(large open squares) and $Z_{\text{sub}} = 0.005 Z_{\odot}$ (large open circles). For comparison, simple stellar populations with metallicities $Z = Z_{\text{maj}}$ and $t = t_{\text{maj}}$ are shown as large open triangles. The dotted (horizontal) lines mark models of constant Z_{sub} .

The perturbation of metal-rich simple stellar populations with metal-poor stars leads to lower $\text{Fe}5335$ line strengths, bluer $B-K$ colors, and stronger Balmer lines. The models with $Z_{\text{sub}} = 0.005 Z_{\odot}$ reach $\text{H}\beta \sim 2$ Å. Observed line strengths (González 1993; Kuntschner & Davies 1998; Mehlert et al. 2000) and colors (Pahre 1998) of elliptical galaxies and lenticular galaxies are shown as small filled and small open symbols, respectively. The present set of composite models match the area that is covered by the majority of the galaxy data. Note that the objects exhibiting $\text{H}\beta > 2$ Å are either classified as lenticular galaxies (small open symbols) or they are intermediate-mass *field* ellipticals (NGC 1700, NGC 3377, NGC 5831, NGC 6702, NGC 7454) or dwarf ellipticals (M32 (NGC 221), NGC 4489) from the González (1993) sample. All *luminous field ellipticals* and all *cluster ellipticals* (Coma, Fornax, and Virgo) in Fig. 3, instead, can be modeled with the old composite populations introduced above. It should be emphasized that the bulk of ellipticals scatter around $\text{H}\beta \sim 1.6$ Å, so that less than 10 per cent contribution from metal-poor stars is required in most cases. The observed range $\text{H}\beta \sim 1.3-2.0$ is then due to a spread in the metallicity and/or the weight of the metal-poor subcomponent. The analysis with simple stellar population models, instead, yields an *age*-spread of $\sim 7-15$ Gyr.

Another striking feature of Fig. 3 is that the positions of the galaxy data relative to each other are the same in the $\text{Fe}5335$ and the $B-K$ diagrams. Moreover, the locations of the model grids relative to the data points in the two panels are in good agreement, so that both the metallic index $\text{Fe}5335$ and the color $B-K$ consistently constrain the metallicity range for the major component to $1-3 Z_{\odot}$. This strongly reinforces the use of $\text{Fe}5335$ as metallicity indicator (see Section 2). In Fig. 4 we show the same data and the same model grid in the $\text{H}\beta$ -Mgb plane. As discussed in Section 2.1, Mgb lines of ellipticals are generally




Fig. 3.— $H\beta$ vs. $Fe5335$ and $B-K$ for elliptical (small filled symbols) and lenticular (small open symbols) galaxies. Data are from González 1993 (circles, Virgo and field, $R_e/8$ aperture, labels indicate NGC numbers), Kuntschner & Davies 1998 (squares, Fornax), and Mehlert et al. 2000 (triangles, Coma). $B-K$ colors are taken from Pahre (1998). The error bars denote average 1σ errors. SSP models for $Z_{SSP} = 0.5, 1, 2, 3 Z_\odot$ are large open triangles, with Z_{SSP} increasing from left to right. Composite models for $Z_{maj} = Z_{SSP}$ and $Z_{sub} = 0.05, 0.005 Z_\odot$ are indicated by large open squares and large open circles. Solid and dotted lines denote fixed Z_{maj} and fixed Z_{sub} , respectively. The metal-poor subpopulation contributes 10 per cent by mass. The ages of the major component and the subcomponent are 14 Gyr and 15 Gyr, respectively.

stronger than any SSP model. This feature is attributed to an enhancement of Mg in luminous ellipticals. As a consequence the metallicities derived from the Mg-indices are substantially higher and provoke severe inconsistencies with observed colors (see also Saglia et al. 2000). As the latter are much less affected by abundance ratio anomalies, the $H\beta$ - $B-K$ plane represents the more reliable constraint on stellar population models.

Note that the $H\beta$ line index additionally suffers from uncertainties in the correction for contamination by emission (González 1993), which can lead to an underestimation of the $H\beta$ line strength. This effect may explain the low values $H\beta \lesssim 1.5 \text{ \AA}$ that fall below the model grid in Fig. 3.

3.2. Constraints from the mid-UV

The fluxes radiated by metal-poor stellar populations are high in the mid-UV (2000–4000 Å), and decrease shortward 2000 Å (Fig. 2). The so-called UV-upturn below 2000 Å observed in elliptical galaxies (Burstein et al. 1988), instead, is most likely due to late evolutionary phases of old metal-rich populations (Greggio & Renzini 1990; Dorman, O’Connell & Rood 1995; Yi, Demarque & Oemler 1998; Greggio & Renzini 1999). The amount of metal-poor stars in elliptical galaxies is therefore tightly constrained by the flux in the mid-UV around 2500 Å.

A model based on the closed-box metallicity distribution is not compatible with the mid-UV flux of the elliptical galaxies

NGC 4649 and NGC 1404 (Bressan, Chiosi & Fagotto 1994) and of the M31 bulge (Worthey, Dorman & Jones 1996), which is referred to as the G-dwarf problem for ellipticals. On the other hand, Lotz, Ferguson & Bohlin (2000) find that the spectra of ellipticals in the mid-UV even require the addition of a metal-poor component. In the following we investigate if the proportions of metal-poor stars required to explain the $H\beta$ line strengths are consistent with the fluxes observed in the mid-UV.

The objects with the strongest $H\beta$ (~ 1.65) that we obtain from cross correlating the samples of González (1993), Kuntschner & Davies (1998), and Mehlert et al. (2000) with the Burstein et al. (1988) sample (‘quiescent objects’) are M31 (NGC 224), NGC 4472, and NGC 3379. We additionally consider NGC 4649, which exhibits a low $H\beta = 1.40$. These objects cover a large range in $1500-V$ colors, NGC 4649 having the strongest UV-upturn. In the following we check if the composite models for these specific galaxies are in agreement with their SEDs. In order to constrain the flux from the metal-poor population in the mid-UV, it is necessary to model the UV-upturn. Following the description by Greggio & Renzini (1990), we include hot ($T \sim 25,000 \text{ K}$) late evolutionary phases in the metal-rich components, such that the observed UV rising branches are reproduced.

The main model parameters for each galaxy are listed in Table 1. Col. (3) specifies the mass fractions (per cent) of the




Fig. 4.— $H\beta$ vs. Mgb . Symbols and linestyles like in Fig. 3.

two components. Note that less than 10 per cent is required in all cases. The metallicities, ages, indices ($H\beta$, $Fe5335$, Mgb), and colors ($B-K$, $U-V$) of the two populations are given in Cols. (4)–(10). The fuel F_{UV} (in M_{\odot}) in late evolutionary phases required to reproduce the observed UV-upturn is given in Col. (11).

In Table 2 we present the relative contributions to $H\beta$ and to the total fluxes at the wavelengths 5550 (V -band), 3600 (U -band), 2500, and 1500 Å. The weight of the metal-poor component increases with decreasing λ , with a maximum contribution in the mid-UV (~ 2500 Å) as the rising branch shortward 2000 Å is produced by the metal-rich component. It is important to emphasize that in the model proposed here, the Balmer line strengths and the UV upturn do not have a common origin: we obtain strong $H\beta$ with *metal-poor* (hot) horizontal branch stars, while blue 1500– V colors are due to extremely hot old *metal-rich* stars. The trend of decreasing $H\beta$ with increasing Mg_2 and decreasing 1500– V (Burstein et al. 1988) is therefore not affected.

In Table 3 we compare the resulting indices and colors of the composed models with the observed values. The indices $H\beta$, $Fe5335$ and the colors $B-K$, $U-V$ are in good agreement with observations.

The fits to the observed SEDs are shown in Fig. 5 as thick solid lines, the dotted lines are the SEDs of the metal-poor and the metal-rich components. The spectra in the UV (open symbols) are from Burstein et al. (1988) and L. Buson (1999, private communication), the spectro-photometry in the optical (thin solid lines) is from D. Hamilton (1999, private communication). In all cases, the observed spectra are well reproduced in the full wavelength range $\lambda = 1200 - 6000$ Å. Thus the amount of metal-poor stars required to reproduce the observed $H\beta$ (see Table 1) is consistent with the fluxes in the mid-UV, the model spectra perfectly match the minimum around 2500 Å. In the following we briefly discuss the individual galaxies.

3.2.1. M31

As shown in Table 1 we reproduce the observables of M31 (for the colors see NGC 4472) with $Z_{\text{sub}} = 0.005 Z_{\odot}$ contributing 3 per cent to the total mass of the population, which cor-

responds to 6 per cent in V , 9 per cent at $H\beta$ and 61 per cent at $\lambda = 2500$ Å. Wu et al. (1980), instead, claim that not more than 50 per cent contribution from a metal-poor component at $\lambda = 2500$ Å is compatible with the mid-UV flux of M31. The reason for this ostensible discrepancy is that the authors fix the metallicity of the metal-rich component to $Z_{\text{maj}} = 0.5 Z_{\odot}$, which is lower than the value adopted here by more than a factor of 3 (see Table 1). As the SED of a more metal-rich population has lower flux in the mid-UV, our composite model requires a larger contribution from the metal-poor component. Using exactly the same prescriptions as these authors, we are consistent with their result. Due to the low metallicity of the metal-rich component, however, the model of Wu et al. (1980) fails in reproducing the other observational constraints of M31: it results in colors that are too blue ($B-K = 3.48$, $U-V \approx 1.21$), Fe indices that are too low ($Fe5335 \approx 2.01$), and Balmer line strengths that are too high ($H\beta = 2.07$).

3.2.2. NGC 4472

The Virgo elliptical NGC 4472 has virtually the same properties as M31, wherefore it can be modeled with the same set of populations. The model given in the second line of Table 1 is an alternative option for both objects assuming a higher contribution from the metal-poor subcomponent and a higher metallicity of the major population. As a consequence, the contribution from the low-metallicity population increases to 70 per cent at 2500 Å and to 12 per cent in V . Wu et al. (1980) derive a lower contribution in V relative to 2500 Å (50 per cent at 2500 Å and 6 per cent in V), because the authors chose the globular cluster M13 (NGC 6205) as a representative of the metal-poor population. As discussed in Section 2.2, this particular globular cluster has an extreme blue horizontal branch morphology and exhibits very high fluxes in the mid-UV (see Fig. 2), which implies a lower V over 2500 Å flux ratio.

3.2.3. NGC 3379

NGC 3379 exhibits equally high $H\beta$, but bluer colors and lower metallic indices. The best fitting model thus has a lower $Z_{\text{maj}} = Z_{\odot}$ and requires only 1 per cent of $Z_{\text{sub}} = 0.005 Z_{\odot}$.

TABLE 1
MODEL PARAMETERS

galaxy	comp.	mass (%)	$Z (Z_{\odot})$	t (Gyr)	$H\beta$	Fe5335	Mgb	$B-K$	$U-V$	$F_{UV} (M_{\odot})$
M31	major	97	1.750	12	1.51	3.20	4.38	4.34	1.73	0.003
	sub	3	0.005	15	3.09	0.53	0.78	2.64	0.53	...
N4472	major	94	2.000	12	1.48	3.32	4.54	4.42	1.81	0.003
	sub	6	0.030	15	2.33	1.02	1.38	2.91	0.72	...
N3379	major	99	1.000	14	1.56	2.82	3.90	4.06	1.59	0.002
	sub	1	0.005	15	3.09	0.53	0.78	2.64	0.53	...
N4649	major	90	1.550	15	1.38	3.21	4.41	4.36	1.73	0.020
	sub	10	0.300	15	1.69	2.08	2.87	3.53	1.15	...

TABLE 2
FLUX CONTRIBUTIONS

galaxy	$H\beta$	5550Å	3600Å	2500Å	1500Å
M31	9	6	17	61	28
N4472	16	12	30	70	16
N3379	4	2	5	33	14
N4649	19	17	27	16	0

NOTE.—The Table gives the relative contributions from the subcomponent in per cent to $H\beta$ (Col. (2)) and to the flux at the wavelengths specified in Cols. (3)–(6).

TABLE 3
RESULTS

galaxy	model	$H\beta$	Fe5335		$B-K$		$U-V$	
		observed	model	observed	model	observed	model	observed
M31	1.65	1.67 ± 0.07	3.04	2.99 ± 0.06	4.26	...	1.61	...
N4472	1.62	1.62 ± 0.06	3.04	2.84 ± 0.08	4.28	4.28	1.59	1.57
N3379	1.62	1.62 ± 0.05	2.77	2.73 ± 0.04	4.05	4.08	1.55	1.52
N4649	1.44	1.40 ± 0.05	3.01	3.01 ± 0.05	4.24	4.26	1.61	1.61

NOTE.—Indices and colors of the models given in Table 1. Index data (1σ errors) are from González (1993), $B-K$ from Pahre (1998), and $U-V$ from Trager (1998).




Fig. 5.— Spectral energy distributions as a function of wavelength. *IUE* data for $\lambda \leq 3300$ Å (circles) are from Burstein et al. 1988 and L. Buson (1999, private communication); the spectro-photometry for $\lambda \geq 3440$ Å (thin solid lines) is from D. Hamilton (1999, private communication). The models from Table 1 are shown as thick solid lines, dotted lines are the SEDs of the metal-poor and the metal-rich components (100 per cent), respectively.

3.2.4. NGC 4649

NGC 4649 has the strongest UV upturn (bluest 1500–V), requiring a factor of ~ 10 more fuel at late evolutionary phases than the other cases (see column (11) in Table 1). Given the low $H\beta$ line strength, the best fit is obtained with a model containing only old metal-rich populations ($Z_{\text{sub}} = 0.3 Z_{\odot}$). This object strongly supports the view that the UV rising branch and $H\beta$ line strength are not produced by the same kind of stars.

3.3. Models for $H\beta > 2$ Å

The model grid in Fig. 3 extends to $H\beta \approx 2$ Å, while some low-mass field ellipticals exhibit stronger Balmer lines. In principle such high $H\beta$ can be obtained by increasing the weight of the low-metallicity component. For instance, the compact dwarf elliptical M32 (NGC 221) with $H\beta = 2.37 \pm 0.12$, $\text{Fe}5335 = 2.54 \pm 0.20$ (Kormendy & Bender 1999) can be modeled by the 14-Gyr old two-component population with $Z_{\text{maj}} = 2-3 Z_{\odot}$ and 20 per cent of $Z_{\text{sub}} = 0.005 Z_{\odot}$. The SED of this

model with such a large fraction of very metal-poor stars, however is not compatible with the fluxes observed in the mid-UV (see also Burstein et al. 1984; Rose & Deng 1999). On the other hand, from a deep *HST* color magnitude diagram, Grillmair et al. (1996) conclude that there is no evidence for an intermediate age population ($t \lesssim 2$ Gyr). It is worth mentioning, however, that M32 is a peculiar object that does not have the properties of the average elliptical galaxy population.

Alternatively to the addition of metal-poor stars, metal-rich populations with blue horizontal branch morphologies could reproduce strong $H\beta$ without invoking young ages. Such hot horizontal branch stars are indeed observed in the metal-rich globular clusters NGC 6388 and NGC 6441 of the galactic bulge (Rich et al. 1997) and in M32 (Brown et al. 2000). A possible mechanism to produce these temperatures is enhanced mass loss along the RGB evolutionary phase. The investigation of this channel to obtain strong $H\beta$ in old metal-rich populations will be the subject of a future paper.

4.1. *Metallicity distributions in ellipticals*

The principle idea of this paper is to enhance the Balmer line strengths of old metal-rich stellar populations with a small fraction of metal-poor stars. The real metallicity distribution of the stellar populations in elliptical galaxies is difficult to assess observationally, as the stars cannot be resolved. The *HST* color magnitude diagram of M32 (Grillmair et al. 1996) and the spectroscopy of K giants in the Bulge (Rich 1988) show that these spheroids contain a tail of low metallicity stars. The closest giant elliptical for which color magnitude diagrams are available is NGC 5128 (Harris, Harris & Poole 1999). Analyzing deep *HST* images in the outer halo of NGC 5128, the authors find a differential metallicity distribution that is well reproduced by two closed-box-like chemical enrichment scenarios implying a large number of extremely metal-poor stars. In particular the differential shape of the distribution implies a formation picture in which a metal-poor and a metal-rich population are formed separately from each other.

This picture gets further support from a number of studies that discover bimodal color and metallicity distributions of globular clusters in at least half of the early-type galaxy population (Zepf & Ashman 1993; Gebhardt & Kissler-Patig 1999). Spectroscopic and photometric investigations indicate that both populations are old (Cohen, Blakeslee & Ryzhov 1998; Kissler-Patig et al. 1998; Kissler-Patig, Forbes & Minniti 1998; Kundu et al. 1999; Puzia et al. 1999). More specifically, Puzia et al. (1999) find that the two populations in NGC 4472 are old and coeval, with metallicities $\sim 0.05 Z_{\odot}$ and $\sim Z_{\odot}$. In the framework of hierarchical structure formation galaxies are built by mergers of smaller objects (e.g., White & Rees 1978; White & Frenk 1991; Kauffmann, White & Guiderdoni 1993). A merger of coeval systems without newly induced star formation would result in a coeval composite population. The accretion of a dwarf galaxy by a larger system (minor merger) qualitatively explains the existence of a metal-poor subpopulation.

There is an additional important effect that is independent of the assumed scheme of galaxy formation. As discussed by Greggio (1997), a sizable fraction of metal-poor stars from the outer parts of a galaxy contaminate the light coming from the center due to projection and orbital mixing (Ciotti, Stiavelli & Braccetti 1995). The authors find that projection effects lower the actual central metallicity by roughly 10 per cent. The metal-poor components in the composite models of M31, NGC 4472 and NGC 3379 (Table 1) dilute the metallicities of the metal-rich populations by 3, 6, and 4 per cent, respectively. Projection effects alone thus may be sufficient to explain the amount of metal-poor stars considered in the present analysis.

4.2. *Evidence against recent star formation*

There are observational indications from the Fundamental Plane (Djorgovski & Davis 1987; Dressler et al. 1987; Bender, Burstein & Faber 1992, 1993; Renzini & Ciotti 1993) and the color-magnitude relation (Bower, Lucey & Ellis 1992) that limit the fraction of the young population to less than 10 per cent. As late star formation leads to low α/Fe ratios (Thomas, Greggio & Bender 1999; Thomas 1999), the addition of a young population provokes inconsistencies with the high α/Fe ratios observed in elliptical galaxies (e.g., Worthey et al. 1992). Also the redshift evolution of colors (Aragón Salamanca et al. 1993), of the Mg- σ relation (Bender, Ziegler & Bruzual 1996), of the Kormendy relation (Ziegler et al. 1999), and of the color gra-

dients in cluster ellipticals (Saglia et al. 2000) leave little space for a significant contribution from a young subcomponent. The addition of an old, metal-poor population as discussed in this paper can reconcile these findings with strong Balmer lines.

Disturbed field ellipticals that have strong H β absorption (Schweizer et al. 1990) and blue optical colors (Schweizer & Seitzer 1992) do not show any signature of recent star formation activity in the infrared colors (Silva & Bothun 1998a, 1998b). As discussed by these authors, recent mergers may not have been accompanied by significant star formation, but metallicity effects are favored to explain the enhanced H β line strengths.

5. SUMMARY

In this paper we show that strong Balmer lines can be produced by composite populations that contain a small fraction of old *metal-poor* stars.

The key to this result is that at low metallicities and old ages our SSP models show a steep increase of H β with decreasing metallicity, in accordance to what is observed in metal-poor globular clusters. We compute composite stellar population models that consist of an underlying old metal-rich population and a small fraction of an old metal-poor subcomponent. Assuming a 10 per cent contribution from metal-poor stars, we construct a grid of models in the H β -Fe5335 and the H β -B-K planes that reproduce H β up to 2 Å and covers most of the elliptical galaxy data. It should be emphasized that the same conclusion holds for all Balmer lines, e.g. H γ and H δ . More specifically, the data of all cluster and luminous field ellipticals can be explained by our models without invoking young ages. The scatter in Balmer line strengths is then caused by a spread in the metallicity and/or the weight of the metal-poor component. Most ellipticals can be modeled with less than 10 per cent of metal-poor stars as the data scatter about H $\beta \sim 1.6$ Å.

We further show that the spectral energy distributions of the composite models that reproduce the indices and colors of representative examples agree well with the observed spectra in the wavelength range 1200–6000 Å. Our models are perfectly compatible with the observed minimum at 2500 Å, that tightly constrains the possible amount of metal-poor stars.

We conclude that the age-metallicity degeneracy for complex systems like elliptical galaxies still remains to be solved. The perturbation with old *metal-poor* stars to obtain strong Balmer lines is alternative to the addition of a *young* metal-rich population (e.g., de Jong & Davies 1997; Kuntschner 2000; Longhetti et al. 2000). The key to discriminate between the two options lies in the evolution of Balmer lines with redshift. A young population of 2–5 Gyr should leave its fingerprints with a significant peak of Balmer line strengths at redshifts $z \sim 0.15$ –0.4, depending on the cosmology, which has not been detected so far (Ziegler & Bender 1997). The model presented here, instead, predicts Balmer lines to become monotonically stronger with redshift.

We are grateful to S. Cassisi, D. Hamilton, D. Mehlert, and L. Buson, for providing electronic versions of models and data. We thank R. Bender and M. Kissler-Patig for the careful reading of the manuscript. For many helpful discussions, we thank N. Drory, L. Greggio, U. Hopp, and R. Saglia. The referee is acknowledged for the very useful comments that improved the first version of the paper. This work was supported by the "Sonderforschungsbereich 375-95 für Astro-Teilchenphysik" of the Deutsche Forschungsgemeinschaft.

REFERENCES

- Aragón Salamanca, A., Ellis, R. S., Couch, W. J., & Carter, D. 1993, *MNRAS*, 262, 764
- Ashman, K. M. & Zepf, S. E. 1998, *Globular Cluster Systems* (Cambridge: Cambridge University Press)
- Bender, R., Burstein, D., & Faber, S. M. 1992, *ApJ*, 399, 462
- . 1993, *ApJ*, 411, 153
- Bender, R., Ziegler, B. L., & Bruzual, G. 1996, *ApJ*, 463, L51
- Bono, G., Caputo, F., Cassisi, S., Castellani, V., & Marconi, M. 1997, *ApJ*, 489, 822
- Bressan, A., Chiosi, C., & Fagotto, F. 1994, *ApJS*, 94, 63
- Brown, T. M., Bowers, C. W., Kimble, R. A., Sweigart, A. V., & Ferguson, H. C. 2000, *ApJ*, in press, astro-ph/9909391
- Burstein, D., Faber, S. M., Gaskell, C. M., & Krumm, N. 1984, *ApJ*, 287, 586
- Burstein, D., Bertola, F., Buson, L. M., Faber, S. M., & Lauer, T. R. 1988, *ApJ*, 328, 440
- Ciotti, L., Stiavelli, M., & Braccisi, A. 1995, *MNRAS*, 276, 961
- Cohen, J. G., Blakeslee, J. P., & Ryzhov, A. 1998, *ApJ*, 496, 808
- Covino, S., Galletti, S., & Pasinetti, L. E. 1995, *A&A*, 303, 79
- de Jong, R. S. & Davies, R. L. 1997, *MNRAS*, 285, L1
- Djorgovski, S. & Davis, M. 1987, *ApJ*, 313, 59
- Dorman, B., O'Connell, R. W., & Rood, R. T. 1995, *ApJ*, 442, 105
- Dressler, A., Lynden-Bell, D., Burstein, D., Davies, R. L., Faber, S. M., Terlevich, R. J., & Wegner, G. 1987, *ApJ*, 313, 42
- Faber, S. M. 1972, *A&A*, 20, 361
- Faber, S. M., Trager, S. C., González, J. J., & Worthey, G. 1995, in *Stellar Populations*, ed. P. C. Van der Kruit & G. Gilmore, IAU Symposium 164 (Dordrecht: Kluwer Academic Publishers), 249
- Fusi Pecci, F., Ferraro, F. R., Bellazzini, M., Djorgovski, S., Piotto, G., & Buonanno, R. 1993, *AJ*, 105, 1145
- Fusi Pecci, F. & Renzini, A. 1975, *A&A*, 39, 413
- Gebhardt, K. & Kissler-Patig, M. 1999, *AJ*, 118, 1526
- Gibson, B. K., Madgwick, D. S., Jones, L. A., Da Costa, G. S., & Norris, J. E. 1999, *AJ*, 118, 1268
- González, J. 1993, Phd thesis, University of California, Santa Cruz
- Greggio, L. 1997, *MNRAS*, 285, 151
- Greggio, L. & Renzini, A. 1990, *ApJ*, 364, 35
- . 1999, *Mem. Soc. Astron. Ital.*, 70, 691
- Grillmair, C. J. et al. 1996, *AJ*, 112, 1975
- Harris, G. L. H., Harris, W. E., & Poole, G. B. 1999, *AJ*, 117, 855
- Harris, W. E. 1996, *AJ*, 112, 1487
- Kauffmann, G., White, S. D. M., & Guiderdoni, B. 1993, *MNRAS*, 264, 201
- Kissler-Patig, M., Forbes, D. A., & Minniti, D. 1998, *MNRAS*, 298, 1123
- Kissler-Patig, M. et al. 1998, *AJ*, 115, 105
- Kormendy, J. & Bender, R. 1999, *ApJ*, 522, 772
- Kundu, A., Whitmore, B. C., Sparks, W. B., Macchetto, F. D., Zepf, S. E., & Ashman, K. M. 1999, *ApJ*, 513, 733
- Kuntschner, H. 2000, *MNRAS*, in press, astro-ph/0001210
- Kuntschner, H. & Davies, R. L. 1998, *MNRAS*, 295, L29
- Lejeune, T., Cuisinier, F., & Buser, R. 1998, *A&AS*, 130, 65
- Longhetti, M., Bressan, A., Chiosi, C., & Rampazzo, R. 2000, *A&A*, 353, 917
- Lotz, J. M., Ferguson, H. C., & Bohlin, R. C. 2000, *ApJ*, in press, astro-ph/9911233
- Maraston, C. 1998, *MNRAS*, 300, 872
- Maraston, C. 2000, in *The chemical evolution of the Milky Way: stars versus clusters*, ed. F. Giovannelli & F. Matteucci
- Maraston, C., Greggio, L., & Thomas, D. 1999, *Ap&SS*, in press, astro-ph/9906088
- Mehlert, D., Saglia, R. P., Bender, R., & Wegner, G. 2000, *A&AS*, 141, 449
- O'Connell, R. W. 1976, *ApJ*, 206, 370
- Pahre, M. A. 1998, Phd thesis, California Institute of Technology
- Puzia, T. H., Kissler-Patig, M., Brodie, J. P., & Huchra, J. P. 1999, *AJ*, 118, 2734
- Renzini, A. 1986, in *Stellar Populations*, ed. C. A. Norman, A. Renzini, & M. Tosi (Cambridge: Cambridge University Press), 213
- Renzini, A. & Buzzoni, A. 1986, in *Spectral evolution of galaxies*, ed. C. Chiosi & A. Renzini (Dordrecht: Reidel), 135
- Renzini, A. & Ciotti, L. 1993, *ApJ*, 416, L49
- Rich, R. M. 1988, *AJ*, 95, 828
- Rich, R. M. et al. 1997, *ApJ*, 484, L25
- Richer, H. B. et al. 1996, *ApJ*, 463, 602
- Rose, J. A. & Deng, S. 1999, *AJ*, 117, 2213
- Saglia, R. P., Maraston, C., Greggio, L., & Bender, R. 2000, *A&A*, submitted
- Schweizer, F. & Seitzer, P. 1992, *AJ*, 104, 1039
- Schweizer, F., Seitzer, P., Faber, S. M., Burstein, D., Dalle Ore, C. M., & González, J. J. 1990, *ApJ*, 364, L33
- Silva, D. R. & Bothun, G. D. 1998a, *AJ*, 116, 85
- . 1998b, *AJ*, 116, 2793
- Thomas, D. 1999, *MNRAS*, 306, 655
- Thomas, D., Greggio, L., & Bender, R. 1999, *MNRAS*, 302, 537
- Trager, S. C. 1998, Phd thesis, University of California, Santa Cruz
- Tripicco, M. J. & Bell, R. A. 1995, *AJ*, 110, 3035
- van Albada, T. S., de Boer, K. S., & Dickens, R. J. 1981, *MNRAS*, 195, 591
- White, S. D. M. & Frenk, C. S. 1991, *ApJ*, 379, 52
- White, S. D. M. & Rees, M. J. 1978, *MNRAS*, 183, 341
- Worthey, G. 1994, *ApJS*, 95, 107
- Worthey, G., Dorman, B., & Jones, L. A. 1996, *AJ*, 112, 948
- Worthey, G., Faber, S. M., & González, J. J. 1992, *ApJ*, 398, 69
- Worthey, G., Faber, S. M., González, J. J., & Burstein, D. 1994, *ApJS*, 94, 687
- Wu, C. C., Faber, S. M., Gallagher, J. S., Peck, M., & Tinsley, B. M. 1980, *ApJ*, 237, 290
- Yi, S., Demarque, P., & Oemler, Jr., A. 1998, *ApJ*, 492, 480
- Zepf, S. E. & Ashman, K. E. 1993, *MNRAS*, 264, 611
- Ziegler, B. L. & Bender, R. 1997, *MNRAS*, 291, 527
- Ziegler, B. L., Saglia, R. P., Bender, R., Belloni, P., Greggio, L., & Seitz, S. 1999, *A&A*, 346, 13
- Zinn, R. & West, M. J. 1984, *ApJS*, 55, 45